



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A NEW BOLETUS FROM MEXICO

W. A. MURRILL

During a recent exploring expedition in southern Mexico, diligent search was made for species of Boletaceae, but only one plant was found, which proved to be undescribed.

Only seven species of this family were previously known from tropical America.*

***Ceriomyces jalapensis* sp. nov.**

Pileus small, convex, circular in outline, 2.2 cm. in diameter, 1 cm. thick; surface isabelline to fulvous, slimy, smooth: context white to faintly roseus, mild to the taste, 2 mm. thick behind; hymenium convex, depressed in the form of a crater about the stipe, tubes pale-greenish, 7 mm. long, mouths large, rounded, 1-2 to a mm., edges thin: spores ellipsoid, deep-ferruginous, distinctly longitudinally striate, copious, $13-15 \times 7-9\mu$: stipe central, slender, tapering upward, concolorous, smooth, glabrous, not conspicuously slimy like the cap, swollen and white at the base, 6 cm. long, 4 mm. thick at the middle.

Type collected near Jalapa, Mexico, at an elevation of 1,700 meters, in humus in a virgin forest, December 12-20, 1909, *W. A. & Edna L. Murrill* 354.

* *Mycologia* 1: 219. 1909.